

Current clinical application of lutetium-177 in solid tumors (Review)

TINGTING NIU^{1,2*}, MI FAN^{1,2*}, BINWEI LIN¹, FENG GAO¹, BANGXIAN TAN² and XIAOBO DU¹

¹Department of Oncology, Mianyang Central Hospital, Mianyang, Sichuan 621000; ²Department of Oncology, Affiliated Hospital of North Sichuan Medical College, Nanchong, Sichuan 637000, P.R. China

Received July 23, 2023; Accepted January 24, 2024

DOI: 10.3892/etm.2024.12514

Abstract. Radionuclide-based therapy represents a novel treatment regimen for tumors. Among these therapies, lutetium-177 (¹⁷⁷Lu) has gained significant attention due to its stability and safety, as well as its ability to emit both γ and β rays, allowing for both imaging with single photon emission computed tomography and tumor treatment. As a result, ¹⁷⁷Lu can be used for both diagnosis and treatment for diseases such as prostatic and gastric cancer. Therefore, based on the available data, the present review provides a brief overview of the clinical applications of ¹⁷⁷Lu-targeted radionuclide therapy in metastatic prostate cancer, neuroendocrine tumors and other types of solid tumors, and highlights the current therapeutic effect, reduction in damage to normal tissues and future research directions, including the development of new nuclides and the application of more nuclides in different tumors. In the future, such treatments could be used in more tumors.

Contents

1. Introduction
2. Basic characteristics of radionuclide ¹⁷⁷Lu
3. Application of ¹⁷⁷Lu in prostate cancer
4. Application of ¹⁷⁷Lu in neuroendocrine tumors (NETs)
5. Clinical application of ¹⁷⁷Lu in other diseases
6. Present problems and future research directions
7. Conclusions

Correspondence to: Professor Xiaobo Du, Department of Oncology, Mianyang Central Hospital, 12 Changjiaxiang, Mianyang, Sichuan 621000, P.R. China
E-mail: duxiaobo2005@126.com

*Contributed equally

Key words: lutetium-177, prostate cancer, neuroendocrine tumor, targeted radionuclide therapy, solid tumors

1. Introduction

The number of total novel cancer cases globally is expected to reach 28.4 million by 2040, which represents a 47% increase compared with 2020 (1). The increase is expected to be higher in developing countries (64-95%) compared with developed countries (32-56%) based on demographic changes (1). On the basis of existing chemotherapy, radiotherapy, immunization, targeting and other therapies, novel diagnosis and treatment methods require investigation. Due to its advantages of integrated diagnosis and treatment, targeted radionuclide therapy has become a new strategy to further improve the prognosis of cancer.

Targeted radionuclide therapy (TRT) involves labeling radionuclides on biomolecules with a specific binding ability, which allows the biomolecule to bind to specific tumor targets, and thus the radionuclide is concentrated at the tumor site (2). The α or β particles emitted by radionuclides can then irradiate the tumor tissues specifically and ionize biological effects (low linear energy transfer), resulting in cell senescence and death within the irradiation range (3). This achieves therapeutic results with minimal effects on the surrounding normal tissues. By combining the advantages of targeted therapy and brachytherapy, TRT provides novel methods for the diagnosis and treatment of tumors. Based on current data, the present review briefly introduced the use of different ¹⁷⁷Lu drugs in treating several tumors, as well as the issues presented by the use of these drugs and future research directions.

2. Basic characteristics of radionuclide ¹⁷⁷Lu

¹⁷⁷Lu belongs to the lanthanide metal group (4) and can emit β -rays with a maximum energy of 0.49 MeV and with a half-life of 6.7 days (5). β -rays are mostly used for treatment of diseases such as prostate cancer, and the energy released by the β -particle rays of ¹⁷⁷Lu has an average range of 670 μ m in soft tissue, significantly reducing the damage caused by radionuclides to nearby healthy cells. In addition, ¹⁷⁷Lu can emit two characteristic γ -rays (208 and 113 keV), which can be used as a signal source for single-photon emission computed tomography or positron emission tomography (PET) (6). These are mostly used for imaging and dosimetric analysis; for example, ¹⁷⁷Lu-prostate-specific membrane antigen (PSMA)-image and treat (I&T) PET/computed tomography (CT) is used for

pre- and post-treatment imaging in patients with metastatic prostate cancer (6,7).

3. Application of ¹⁷⁷Lu in prostate cancer

The latest global cancer data results released by the World Health Organization shows that the number of new cases of prostate cancer in 2020 was 1.41 million, ranking it second most common amongst all new tumors and the number of prostate cancer-associated mortalities in men was 380,000, making it the fifth most fatal cancer in men (1). In recent years, the prevalence of metastatic castration-resistant prostate cancer (mCRPC) has been increasing, and its prognosis is generally poor (8). The expression of PSMA in prostate cancer is higher compared with that in other tissues, including prostatic epithelial cells, the small intestine and salivary glands (9), and is 1,000 times higher compared with the lowest expression level in the kidney and small intestine (10). Therefore, precise targeted therapy with PSMA has emerged as a novel therapeutic method, resulting in a decline in prostate specific antigen (PSA) levels and an increase in overall survival (11,12).

Prostate cancer and ¹⁷⁷Lu-PSMA-I&T. 1,4,7,10-tetraazacyclododecane-1-(glutaric acid)-4,7,10-triacetic acid (DOTAGA) is a small PSMA inhibitor molecule labeled with ¹⁷⁷Lu known as ¹⁷⁷Lu-PSMA-I&T. In 2015, internal radiation therapy using ¹⁷⁷Lu-PSMA-I&T was reported to be effective and safe in two patients with metastatic prostate cancer, with no detectable side effects (13). In a study by Barna *et al* (14), ¹⁷⁷Lu-PSMA-I&T was used to treat mCRPC with a mean injection activity of 7,416±218 MBq. Follow-up imaging using Ga-PSMA PET/CT was used to determine individual tumor molecular volume. The volume of 63 individual tumors in bone, lymph nodes and liver tissues were observed to decrease by 32.3, 84.7 and 72.9% on average, respectively (14). In a trial of 49 patients with mCRPC who received at least three cycles of ¹⁷⁷Lu-PSMA-I&T (6.0 GBq), no grade III/IV adverse events were reported, indicating low nephrotoxicity or hematotoxicity, according to the Common Terminology Criteria for Adverse Events (CTCAE v5.0) (15).

Another prospective single-arm trial included 40 patients diagnosed with mCRPC who received 86 cycles of ¹⁷⁷Lu-PSMA-I&T at a dose of 3.70-14.43 GBq per cycle from December 2019 to September 2021, with a median follow-up of 8 months (16). The findings revealed that six patients (15%) developed mild reversible dry mouth, and 28 patients (70%) developed grade 1-4 bone marrow dysfunction (anemia, thrombocytopenia and leukopenia) during follow-up (toxicities were assessed following the CTCAE v5.0). Serum PSA is the most significant marker for evaluating therapeutic biochemical response. PSA levels were obtained every 4 weeks before and after treatment. A reduction of ≥30% from baseline is considered a partial response, a >25% increase in PSA above baseline is defined as disease progression, and PSA levels with changes between <-30% and <+25% are considered disease stabilization. PSA levels were assessed after treatment and accompanied by partial remission in 25 patients (62.5%), stable disease in five patients (12.5%) and progression in ten patients (25%). Trials have shown that ¹⁷⁷Lu-PSMA-I&T can

achieve significant PSA reduction and tumor remission in patients with mCRPC (16,17).

To evaluate the predictive effect after treatment, a trial involving 301 patients demonstrated that the extent of bone metastases and their changes were potential markers for predicting treatment outcomes in patients with mCRPC (18). A previous single-arm phase I study of ¹⁷⁷Lu-PSMA-I&T neoadjuvant therapy in high-risk prostate cancer before radical prostatectomy included 14 patients with high-critical-limited-stage prostate cancer, defined as PSA >20 ng/ml, biopsy Gleason score ≥8, or clinical T stage ≥3a, who tested positive for PSMA between December 2019 and December 2021 (19). Furthermore, two or three doses of ¹⁷⁷Lu-PSMA-I&T (7.4 GBq) were administered at 2-week intervals, and surgery (including lymph node dissection) was performed 4 weeks after the final dose. The main adverse reactions of this treatment were the incidence of perioperative complications and organ functional toxicity, including bleeding, infection, pneumonia and pulmonary embolism. After two doses, the PSA level was reduced by 17%, and after three doses it was reduced by 34%. No severe intraoperative complications were observed in 13 patients (one patient did not undergo surgery due to heart problems), while four (30%) patients developed postoperative complications (including pneumonia, pulmonary embolism, urinary leakage and urinary tract infection) (19).

Preliminary studies have demonstrated that neoadjuvant therapy with ¹⁷⁷Lu-PSMA-I&T followed by surgery is safe; however, further data should be obtained from long-term follow-up (19). These findings suggest that patients treated with ¹⁷⁷Lu-PSMA-I&T show decreased PSA levels and fewer serious adverse events, contributing to tumor remission (16-18).

Prostate cancer and ¹⁷⁷Lu-PSMA-617. ¹⁷⁷Lu-PSMA-617 is a small molecule inhibitor labeled with ¹⁷⁷Lu using 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (DOTA) as a chelating agent (20,21). In March, 2022, the U.S. Food and Drug Administration (FDA) approved the administration of ¹⁷⁷Lu-PSMA-617 for the treatment of adult patients with mCRPC, making it the first targeted radioligand therapy to be approved by the FDA (22).

In a meta-analysis, Kim and Kim (23) included 10 studies involving 455 patients with mCRPC. The analysis showed that ¹⁷⁷Lu-PSMA-617 therapy resulted in a combined PSA reduction rate of 68.00% [95% confidence interval (CI), 63.55-72.22%] and with PSA levels reduced by >50% in 34.45% of patients (95% CI, 30.14-38.97%). In phase II, single-arm, single-center trials by Violet *et al* (24) and Hofman *et al* (25), ¹⁷⁷Lu-PSMA-617 radionuclide therapy had a high response rate, low toxicity and decreased pain in patients with metastatic castration-tolerant prostate cancer who progressed following conventional therapy (including taxane-based chemotherapy and second-generation antiandrogen therapy). The FDA authorized ¹⁷⁷Lu-PSMA-617 as a prostate cancer treatment in March 2022 based on a phase III experiment conducted by Sartor *et al* (26).

An international, open-label, Phase III clinical trial involving 84 centers (52 centers in North America and 32 in Europe) was conducted (26). Patients were randomly assigned in a 2:1 ratio to receive either ¹⁷⁷Lu-PSMA-617 (injection of 7.4 GBq once every 6 weeks, lasting 4-6 cycles) plus standard

protocol-approved treatment (^{177}Lu -PSMA-617 group) or standard treatment only (e.g., abiraterone and enzalutamide) (control group). Among the 831 patients with metastatic castration-tolerant prostate cancer treated with at least one androgen receptor pathway inhibitor and one or two taxane receptor inhibitors, 581 were included in the analysis set. ^{177}Lu -PSMA-617 + standard therapy significantly extended imaging-based progression-free survival (PFS) time compared with standard therapy (median, 8.7 vs. 3.4 months; hazard ratio for progression or death, 0.40; 99.2% CI, 0.29-0.57; $P < 0.001$) and OS (median, 15.3 vs. 11.3 months; hazard ratio for death, 0.62; 95% CI, 0.52-0.74; $P < 0.001$). The most frequent adverse reactions in the ^{177}Lu -PSMA-617 group were fatigue, dry mouth and nausea; however, these events were usually grade 1 or 2 (Response Evaluation Criteria in Solid Tumors, version 1.1) (27) and had little impact on the quality of life.

^{177}Lu -PSMA-I&T vs. ^{177}Lu -PSMA-617. The distinction between the two PSMA-targeting medications, ^{177}Lu -PSMA-617 and ^{177}Lu -PSMA-I&T, lies in the chelating agents they use. A study with 110 patients with mCRPC from two locations (University Hospital Würzburg and University Hospital Bonn), including 55 who received ^{177}Lu -PSMA-I&T and 55 who received ^{177}Lu -PSMA-617, revealed no significant differences in the harmful effects of the two at a dose of roughly 6.0 GBq per 8 weeks. Survival with ^{177}Lu -PSMA-I&T and ^{177}Lu -PSMA-617 was comparable with a median OS time of 12.0 vs. 13.0 months, respectively, with no serious grade III/IV toxicity (28). In another study that assessed the safety, biological distribution and dosiology in 138 patients, 51 individuals were administered ^{177}Lu -PSMA-I&T at 6.1 ± 1.0 GBq and 87 patients received ^{177}Lu -PSMA 617 at 6.5 ± 1.1 GBq (one injection). The mean dose of ^{177}Lu -PSMA-617 was higher compared with ^{177}Lu -PSMA-I&T (0.04 vs. 0.03 Gy/GBq), and the systemic half-life of ^{177}Lu -PSMA-I&T (35 h) was shorter compared with that of ^{177}Lu -PSMA-617 (42 h). Of all the healthy organs, the lacrimal glands had the highest mean absorbed tumor dose of ^{177}Lu -PSMA-I&T and ^{177}Lu -PSMA-617 (5.8 vs. 5.9 Gy/GBq), but patients tolerated the therapy without any acute side effects (29). A study that further evaluated the difference in efficacy between ^{177}Lu -PSMA-617 plus standard therapy (hormone therapy, bisphosphonates and radiotherapy) and ^{177}Lu -PSMA-617 alone showed that combined therapy extended the duration of pain exacerbation compared with ^{177}Lu -PSMA-617 alone (30). It is hoped that future studies can include the combined use of nuclides with other drugs (such as targeted drugs and chemotherapy drugs), and it is hoped that new progress and breakthroughs can be made in the treatment of mCRPC through combination therapy.

Prostate cancer and ^{177}Lu -J591. J591 is a deimmunizing monoclonal antibody that binds to PSMA and has internalized properties (internalization of a putative ligand) (31,32). Participants received 75 mg/m docetaxel once every 21 days for 2 cycles, with two progressive increments of 'Lu-1591' at cycle 3 (1.48 GBq/m², up to 2.96 GBq/m²). Docetaxel was administered in cycle 3 (half of the Lu-J591 dose 2 to 3 days before docetaxel and the other half of the dose 2 weeks after docetaxel), with the fourth cycle of docetaxel given 6-9 weeks after the third cycle. No dose-limiting toxicity was observed

at any of the tested dose levels. Grade 4 neutropenia without fever was found in eight patients (53.5%). Subsequently, two patients (13.3%) developed thrombocytopenia, and no grade ≥ 3 non-hematological toxicity was observed. A PSA decrease of $>50\%$ after treatment with ^{177}Lu -J591 was seen in 11 patients (73.3%). The results suggest that a single ^{177}Lu -J591 fractionated course in combination with docetaxel is a viable option for patients with mCRPC (33). A review of the various phase I and II trials of ^{177}Lu -J591 in metastatic prostate cancer showed improved OS, and in almost all cases, myelosuppression is tolerable and reversible (31). While ^{177}Lu -J591 may be considered a potential treatment option, the effectiveness of its combination with docetaxel still requires additional research and clinical trials to establish its efficacy and safety.

Prostate cancer and [^{177}Lu] Ludotadipep. Ludotadipep is a new PSMA inhibitor tagged with ^{177}Lu , which contains a 4-iodophenyl butanoic group that can bind to albumin and prolong circulation time and boost absorption in tumors. A prospective study from Korea evaluated the efficacy of ^{177}Lu Ludotadipep in patients with ^{18}F -PSMA-PET/CT-positive mCRPC. From November 2020 to March 2022, a total of 30 patients were enrolled for single dose of ^{177}Lu Ludotadipep radiopharmaceutical therapy (34). Patients were divided into five groups ($n=6$) and given an increasing dose of 1.9, 2.8, 3.7, 4.6 and 5.6 GBq for each group. In this study, 29 patients who received ^{177}Lu Ludotadipep (one patient found to deviate from the inclusion criteria after enrollment), 36 treatment-emergent adverse events (58.6%) and four adverse drug reactions (10.3%) were observed. Overall 16 (66.7%) of the 24 participants with complete 12-week follow-up data showed a reduction in PSA levels, and nine (37.5%) of those subjects showed a PSA decline of $\geq 50\%$. At the 12th week after receiving a single dosage of ^{177}Lu Ludotadipep, five of the 24 patients (20.8%) displayed disease progression (a $\geq 25\%$ increase in PSA levels from the baseline), suggesting that ^{177}Lu Ludotadipep may be a promising new treatment for mCRPC (34).

4. Application of ^{177}Lu in neuroendocrine tumors (NETs)

NETs are rare, heterogeneous tumors that originate from cells of the diffuse endocrine system (35). Approximately two-thirds of NETs occur in the gastrointestinal and pancreatic systems, including the stomach, small intestine, colon, appendix, rectum and pancreas. The most common NET subtype is the gastrointestinal pancreatic NET (GEP-NET) (36). An analysis of United States cancer data showed that the incidence of gastrointestinal neuroendocrine tumors has continuously increased each decade from 1977 to 2016 (37). Currently, surgery is the primary therapeutic method for GEP-NETs because most GEP-NETs are inert. In addition, severe cases are treated with radiochemotherapy, molecularly targeted medicine.

Most pancreatic and GEP-NETs express somatostatin receptor (SSTR)2 and 5, these receptors can be chelated to a β -emitting radioisotope ^{177}Lu for therapy. The new approach for treating advanced NETs is peptide receptor radionuclide therapy (38).

Somatostatin is a peptide with a potent and broad antisecretory action. SSTRs which belong to the G protein-coupled receptor family, are widely distributed in various tissues of

the body and are classified into five subtypes (SSTR1-5) (39). SSTR2 is associated with gastrointestinal neuroendocrine system (40).

NETs and ¹⁷⁷Lu-1, 4, 7, 10-tetraazacyclododecane-1, 4, 7, 10-tetraacetic acid-D-phenylalanine 1-tyrosine 3-threonine 8-octreotide (¹⁷⁷Lu-DOTA-TATE) also known as DOTA-octreotate. ¹⁷⁷Lu-DOTA-TATE is a labeled precursor of DOTA-modified octreotide. Octreotide is a somatostatin analogue that is used to control NET progression (41). A ¹⁷⁷Lu-labeled SSTR agonist has been used to treat somatostatin receptor positive GEP-NETs (42). The FDA approved the first radiopharmaceutical for the treatment of GET on January 26, 2018.

An open, randomized, phase III clinical trial (NETTER-1) enrolled 231 patients with locally advanced or metastatic, well-differentiated, somatostatin receptor-positive midgut neuroendocrine tumors. Patients were randomly assigned (1:1) to receive intravenous ¹⁷⁷Lu-DOTA-TATE 7.4 GBq + intramuscular long-acting octreotide (30 mg every 8 weeks) (experimental group) or high-dose long-acting octreotide (60 mg every 4 weeks) (control group). According to the NETTER-1 trial, which was reported on July 24, 2016, the estimated rate of PFS at 20 months was 65.2% (95% CI, 50.0-76.8%) in the experimental group and 10.8% (95% CI, 3.5-23.0%) in the control group. The response rate (complete plus partial) in the experimental group was 18%, whereas it was only 3% in the control group (P<0.001). Preliminary findings show that using ¹⁷⁷Lu-DOTA-TATE markedly increases the PFS (43). In a 2021 follow-up analysis the median OS for the experimental group was 48.0 months (95% CI, 37.4-55.2 months), while the control group OS was 36.3 months (95% CI, 25.9-51.7 months). The OS of the experimental group did not improve significantly compared with that of the control group, and there was no statistically significant difference in OS. However, in absolute terms, there was a difference of 11.7 months in the median OS between ¹⁷⁷Lu-DOTA-TATE and octreotide alone groups. In the experimental group, 111 patients (3%) experienced serious adverse events related to treatment that were grade 3 or worse. During the 100 months of follow-up, 2% of patients developed myelodysplastic syndrome, yet no other experimental group patients experienced renal impairment that was grade 3 or worse. Although the OS was not statistically significant, the results showed that the 11.7 monthly difference in the median OS between ¹⁷⁷Lu-DOTA-TATE treatment and high-dose long-acting octreotide treatment alone may be considered clinically relevant (44).

A study compared PFS and OS in ¹⁷⁷Lu-DOTA-TATE and patients with advanced and unresectable gastrointestinal neuroendocrine tumors treated with everolimus and sunitinib (45). In comparison to everolimus, sunitinib and best supportive care, the results of the primary analysis demonstrated that ¹⁷⁷Lu-DOTA-TATE may be a more effective therapy choice. In a study evaluating the efficacy and safety of ¹⁷⁷Lu-DOTA-TATE in 30 patients with NET and extensive bone metastases, radiological evaluation at the end of treatment showed partial response in five patients, stable disease in 20 patients, and progressive radiological disease in three patients. Clinical progress was observed in another two patients (46).

NETs and ¹⁷⁷Lu-DOTA-JR11 (¹⁷⁷Lu-OPS201). ¹⁷⁷Lu-OPS201 is a novel somatostatin antagonist with high affinity for SSTR2 (47). *In situ* testing in a SSTR2-positive neuroendocrine model in mice reveals that ¹⁷⁷Lu-OPS201 causes a greater decrease in living tumor tissue, a significant delay in tumor growth and increased toxicity compared with ¹⁷⁷Lu-DOTATOC (an SSTR agonist that primarily targets SSTR2). Likewise, the use of the ¹⁷⁷Lu-OPS201 has been shown to increase tumor uptake of agonist ¹⁷⁷Lu-DOTA-TATE *in vitro* (47); however, it has not been validated at clinical trial stage. According to a recent *in vitro* study, ¹⁷⁷Lu-OPS201 has a high affinity, and ¹⁷⁷Lu-OPS201 has at least four times more receptor-binding sites compared with ¹⁷⁷Lu-DOTA-TATE. In conclusion, ¹⁷⁷Lu-OPS201 has demonstrated faster binding, slower dissociation and a longer cell retention period compared with ¹⁷⁷Lu-DOTA-TATE (48).

5. Clinical application of ¹⁷⁷Lu in other diseases

Fibroblast activating protein (FAP) overexpression in cancer and ¹⁷⁷Lu-FAP-2286. FAP is expressed in a number of malignancies. ¹⁷⁷Lu-FAP-2286 is a FAP-binding cyclic peptide consisting of seven amino acids, of which two cysteine residues pass through an aromatic partial ring that is linked to the DOTA chelating agent (49). FAP-2287 (a murine surrogate for FAP-2286) is well targeted for ¹⁷⁷Lu targeted radionuclide therapy, rapidly accumulates in tissues and persists in tumors for a long time (50).

A trial has been performed that involved 11 patients with advanced pancreatic, breast, rectal and ovarian cancers receiving ¹⁷⁷Lu-FAP-2286. The results showed that the dosage of administration of ¹⁷⁷Lu-FAP-2286 (5.8±2.0 GBq; range, 2.4-9.9 GBq) was well tolerated, and no adverse or clinically detectable pharmacological effects were found or reported in any patient. The systemic effective dose was 0.07±0.02 Gy/GBq (range, 0.04-0.1 Gy/GBq). The mean absorbed doses in kidney and red bone marrow were 1.0±0.6 Gy/GBq (range, 0.4-2.0 Gy/GBq) and 0.05±0.02 Gy/GBq (range, 0.03-0.09 Gy/GBq), respectively. No grade 4 adverse events were observed; however, grade 3 adverse reactions occurred in three patients, including one pancytopenia, one leukopenia and one pain response. ¹⁷⁷Lu-FAP-2286 is widely used in adenocarcinoma and is well tolerated with few side effects (51). These results indicate that ¹⁷⁷Lu-FAP-2286 has encouraging clinical data and deserves further exploration.

Meningioma and ¹⁷⁷Lu-DOTA-TATE. At present, there is no evidence-based systemic treatment for patients with progressive meningioma who are unable to receive surgery or external radiotherapy. External radiation therapy was given to 15 patients with meningioma, all of whom had received radiotherapy and 14 of whom had surgery and then received ¹⁷⁷Lu-DOTA-TATE (7.5-29.6 GBq). ¹⁷⁷Lu-DOTA-TATE was administered with a maximum activity of 7.4 GBq per cycle with a maximum of 4 cycles. In this cohort, the interval between the cycles was a median of 9 weeks (range, 6-14 weeks). Subsequently, six patients (40%) were stable following treatment. The median PFS of the whole cohort was 7.8 months, with a 6-month PFS rate of 60%. The median OS was 13.6 months, with a 12-month OS rate of 60%. Prior to therapy, their average monthly tumor

growth rate (TGR) was 4.6% for the surface and 14.8% for the volume. The surface and volume were scanned and determined by magnetic resonance imaging. The TGR decreased to 3.1% in surface (P=0.016) and 5.0% in volume (P=0.013) per month after treatment. The results suggest that ¹⁷⁷Lu-DOTA-TATE can control tumor growth (52).

Metastatic salivary gland cancer and ¹⁷⁷Lu-PSMA-617. PSMA is expressed on tumor cells or in the tumor neovasculature in salivary gland carcinoma, particularly in certain subtypes such as salivary gland and salivary duct carcinomas (53,54). There have been cases of salivary gland malignancies that show significant uptake in ⁶⁸Ga-PSMA-11 PET-CT (55-57). Therefore, in view of the encouraging results of patients treated for CRPC, the potential use of ¹⁷⁷Lu-PSMA-617 to treat metastatic salivary gland cancer has been assessed.

In a retrospective study by Klein Nulent *et al* (58), six patients were treated with ¹⁷⁷Lu-PSMA-617. This cohort included four adenoid cystic carcinomas, one adenocarcinoma (not otherwise specified) and one acinic cell carcinoma. A total of four patients reported instant reduction of tumor-related symptoms; the most common improvement was a reduction in pain, followed by a reduction in fatigue. Two individuals demonstrated a radiological response, indicating either stable disease or a partial remission. All treatment-related clinical and hematological adverse events were graded using the CTCAE standard version 5.0. This study demonstrated that palliative ¹⁷⁷Lu-PSMA-617 for salivary gland cancer is safe and generally well tolerated.

A 56-year-old man with progressive metastatic salivary cancer who received treatment every 6 weeks and was first evaluated after four courses (cumulative activity of 24.3 GBq ¹⁷⁷Lu-PSSMA-617) showed stable disease on imaging according to Positron Emission tomography Response Criteria In Solid Tumor (PERCIST) criteria (59).

Metastatic thyroid cancer and ¹⁷⁷Lu-DOTA-TATE/¹⁷⁷Lu-P SMA-617I. A number of studies have shown that PSMA is expressed in thyroid cancer, and significant uptake can be observed with ⁶⁸Ga-PSMA-11 PET-CT (60-63). Therefore, ¹⁷⁷Lu-PSMA-617 has been used to treat metastatic thyroid cancer. Assadi and Ahmadzadehfar (64) treated a patient with progressing thyroid cancer and neck and lung metastases with ¹⁷⁷Lu-PSMA. The patient previously received 25.9 GBq ¹³¹Iodine and sorafenib for 6 months, and radioligand therapy targeting the somatostatin receptor using ¹⁷⁷Lu-DOTA-TATE. As the patient exhibited persistent disease progression and severe breathing difficulties, the patient was then treated with 7.4 GBq ¹⁷⁷Lu-PSMA-617. With whole-body SPE-CT imaging, there was a higher level of whole-body imaging with ¹⁷⁷Lu-PSMA-617 compared with receiving ¹⁷⁷Lu-DOTA-TATE, suggesting that treatment with ¹⁷⁷Lu-PSMA-617 is potentially more effective compared with ¹⁷⁷Lu-DOTA-TATE for the treatment of this patient. However, the patient died unexpectedly of cardiac arrest two weeks after treatment.

In an additional study, two patients received ¹⁷⁷Lu-PSMA-617 therapy for thyroid carcinoma. One patient showed disease progression on imaging 1 month later, the other patient showed partial, temporary response of lung and

Table I. Recruiting clinical trials of ¹⁷⁷Lu targeted radionuclide therapy for tumors expressing SSTR and PSMA (except neuroendocrine tumor and prostate cancer) in the past 5 years.

NCT	Starting date	Phase	Target	Vector	Isotope	Application	Administration Route	Primary outcome
NCT04529044	2022-09-01	II	SSTR	Dotatate	Lu-177	Recurrent breast cancer	I.V	ORR
NCT05198479	2023-01	II	SSTR	DOTA0-Tyr3-Octreotate	Lu-177	Metastatic NPC	I.V	PFS
NCT04903899	2021-05-19	II	SSTR	DOTATATE	Lu-177	Neuroblastoma	I.V	Assess the response of treatment.
NCT05644080	2023-03-28	N/A	PSMA	PSMA I&T	Lu-177/Ga-68	Glioma	I.V	AEs/efficacy
NCT05918302	2023-07	III	SSTR	Edotreotide	Lu-177	Neuroendocrine tumors	I.V	PFS
NCT05109728	2022-05-10	I	SSTR	DOTA-TATE	Lu-177	Glioblastoma	I.V	DLTs
NCT03971461	2019-05-15	II	SSTR	Lutathera	Lu-177	Meningioma	I.V	PFS
NCT05214820	2022-01-17	II	PSMA	⁶⁸ Ga-PSMA	Ga-68	Endoradiotherapy	I.V	Tumor uptake

SSTR2, somatostatin receptor subtype 2; PSMA, prostate-specific membrane antigen; NPC, metastatic nasopharyngeal cancer; ORR, overall response rate; PFS, progression free survival; DLTs, number of participants with dose limiting toxicities; AEs, adverse events; NCT, national clinical trial; I.V, intravenous; PFS, progression-free survival; N/A, not applicable; I&T, image and treat.

Table II. New targets of clinical trials of ¹⁷⁷Lu targeted radionuclide therapy for solid tumors in the past 5 years.

NCT	Starting date	Phase	Target	Vector	Isotope	Application	Administration route	Primary outcome
NCT05623891	2022-12	I	ED-B	B5-IgG4	Lu-177	Solid Tumors	I.V	Tissue distribution/ Dosimetry/AEs
NCT05815394	2023-03-03	I	NY108	DOTA	Lu-177	Prostate cancer	I.V	Tissue distribution/AEs
NCT04786847	2022-01-30	I	PSMA	TLX591	Lu-177	MPC	I.V	AEs
NCT05723640	2023-05-10	I	FAP	LNC1004	Lu-177	Solid tumor	I.V	DLTs/safety/MDT
NCT05410821	2022-06-15	I	FAP	DOTA-EB-FAPI	Lu-177	Thyroid cancer	I.V	Safety/tolerability/MDT
NCT05603559	2023-01-01	I	PSMA	P17-087/088	Lu-177	mCRPC	I.V	Dosimetry/AEs
NCT05013086	2021-10-01	I	Integrin $\alpha v \beta 3$	AB-3PRGD2	Lu-177	NSCLC	I.V	Standardized uptake value
NCT05130255	2022-11-17	I	GD2	DOTA	Lu-177	Solid tumors	I.V	DLTs
NCT04997317	2021-04-21	I	SSTR2	satoreotide	Lu-177	Meningiomas	I.V	Therapeutic index/safety
NCT03872778	2019-07-24	I/II	GRPR	NeoB	Lu-177	Solid tumor	I.V	DLTs/MDT
NCT04665947	2020-12-18	I	ABM-5G	DOTA	Lu-177	Pancreatic cancer	I.V	DLTs/RP2D
NCT05706129	2023-03-14	I/II	DPI-4452	DOTA	Lu-177/Ga-68	Solid tumors	I.V	TEAEs/DLTs/ORR
NCT05868174	2023-05-23	I	CAIX	Pepeortib	Lu-177	Solid tumors	I.V	DLTs/safety
NCT05432193	2022-07-13	I	FAP	PNT6555	Lu-177	Solid tumors	I.V	AEs
NCT04711135	2022-08-31	II	SSTR	Lutetium	Lu-177	GEP-NETs	I.V	Absorbed radiation doses/AEs
NCT04647526	2021-02-25	III	PSMA	PNT2002	Lu-177	mCRPC	I.V	rPFS
NCT04469127	2023-01-30	I/II	$\alpha v \beta 3$	DOTA	Lu-177	Breast cancer	I.V	Safety and tolerability
NCT05533242	2023-06	I	6A10	DOTA	Lu-177	Glioblastoma	I.V	MDT/safety
NCT05178693	2022-04-25	I	SSTR	ASTX727	Lu-177	NETs	I.V	To determine whether pre-treatment with ASTX727 results in re-expression of SSTR2

PSMA, prostate-specific membrane antigen; FAP, fibroblast activation protein; SSTR2, somatostatin receptor subtype 2; GRPR, gastrin-releasing peptide receptor; CAIX, carbonic anhydrase IX; MPC, metastatic prostate cancer; mCRPC, metastatic castration-resistant prostate cancer; NSCLC, non-small cell lung cancer; GEP-NETs, gastroenteropancreatic neuroendocrine tumors; I.V, intravenous; AEs, adverse events; DLTs, number of participants with dose limiting toxicities; MTD, maximum tolerated dose; RP2D, recommended phase 2 dose; TEAEs, treatment emergent adverse events; ORR, overall response rate; rPFS, radiographic progression-free survival; NCT, national clinical trial.

Table III. Ongoing clinical trials of ¹⁷⁷Lu targeted radionuclide therapy combined with other solid tumor treatments in the past 5 years.

NCT	Starting date	Phase	Target	Combination treatment	Application	Primary outcome
NCT05340374	2022-07-14	I/II	PSMA	Cabazitaxel + ¹⁷⁷ Lu-PSMA-617	mCRPC	DLTs/MTD/RP2D
NCT04525638	2020-06-29	II	SSTR	Nivolumab + ¹⁷⁷ Lu-DOTATATE	NENS	ORR
NCT05766371	2023-06-30	II	PSMA	Pembrolizumab + ¹⁷⁷ Lu-PSMA-617	CRPC	rPFS
NCT03874884	2019-07-09	I	PSMA	Olaparib + ¹⁷⁷ Lu-PSMA-617	mCRPC	DLTs/MTD/RP2D
NCT05150236	2022-04-29	II	PSMA	Ipilimumab + Nivolumab + ¹⁷⁷ Lu-PSMA-617	mCRPC	PSA-PFS
NCT04886986	2021-06-30	I/II	PSMA	²²⁵ Ac-J591 + ¹⁷⁷ Lu-PSMA-I&T	mCRPC	DLTs/MTD
NCT05613894	2023-06-26	Ib	PSMA	Cabozantinib + ¹⁷⁷ Lu-PSMA-617	mCRPC	DLTs
NCT05239533	2022-02-16	II	Anti-carbonic anhydrase IX monoclonal antibody	Nivolumab + ¹⁷⁷ Lu-girentuximab	Kidney cancer	MTD/ORR
NCT05868174	2023-05-23	I	CAIX	Peposertib + ¹⁷⁷ Lu-TLX250	Solid tumors	DLT/safety
NCT04720157	2021-06-09	III	PSMA	SoC + ¹⁷⁷ Lu-PSMA-617	mHSPC	rPFS
NCT05663710	2023-06-30	II	SSTR	Cabozantinib + Nivolumab	RCC	Incidence of PFS
NCT04261855	2020-10-08	I/II	SSTR	Avelumab + ¹⁷⁷ Lu-DOTATATE	MCC	PFS
NCT05383079	2022-09-13	I/II	PSMA	²²³ Ra + ¹⁷⁷ Lu-PSMA-I&T	mCRPC	DLTs/MTD/RP2D
NCT05724108	2023-03-15	II	SSTR	Triapine + ¹⁷⁷ Lu-DOTATATE	NENS	ORR
NCT05249114	2022-12-28	I	SSTR	Cabozantinib + ¹⁷⁷ Lu-DOTATATE	NENS	MTD PFS
NCT05687123	2023-04-28	I	SSTR	Sunitinib Malate + ¹⁷⁷ Lu-DOTATATE	Pancreatic NENS	AEs
NCT04750954	2021-06-04	I	SSTR	Peposertib + ¹⁷⁷ Lu-DOTATATE	Pancreatic NENS	RP2D
NCT05682443	2023-07-01	II	PSMA	ONC-392 + ¹⁷⁷ Lu-PSMA-617	mCRPC	rPFS
NCT05247905	2022-03-16	II	SSTR	Capecitabine + Temozolomide	Pancreatic NENS	PFS
NCT04086485	2022-10-03	I/II	SSTR	Olaparib + ¹⁷⁷ Lu-DOTATATE	GEP-NET	MTD/ORR
NCT04614766	2022-09-30	I/II	SSTR	Lutathera + Azedra	SPORE-3	MTD/ORR

PSMA, prostate-specific membrane antigen; CAIX, carbonic anhydrase IX; SSTR, somatostatin receptor subtype; SoC, standard of care; mCRPC, metastatic castration resistant prostate cancer; OPC, oligometastatic prostate cancer; NENS, neuroendocrine neoplasms; CRPC, castration resistant prostate cancer; mHSPC, metastatic hormone sensitive prostate cancer; RCC, renal cell carcinoma; MCC, Merkel cell carcinoma; GEP-NET, gastroenteropancreatic neuroendocrine tumor; SPORE-3, mid-gut neuroendocrine tumor; SPORE-3, number of participants with dose limiting toxicities; MTD, maximum tolerated dose; RP2D, recommended Phase 2 dose; PSA PFS, prostate-specific antigen progression free survival at 1 year; bPFS, biochemical progression free survival; ORR, overall response rate; rPFS, radiographic progression-free survival; AEs, adverse events.

liver metastases (65). Side effects were not mentioned in either of these two case reports of thyroid carcinoma.

Advanced renal cell cancer and ¹⁷⁷Lu-cG250. Previous studies have shown that carbonic anhydrase IX is almost universally expressed (>90%) in metastatic clear cell renal cell carcinoma (ccRCC), and its expression in healthy tissues has been extensively evaluated, but is limited to gastrointestinal mucosa and gastrointestinal associated structures, with much lower expression levels compared with those in ccRCC (66,67). Thus, ¹⁷⁷Lu-girentuximab (cG250) was used to treat ccRCC. Diagnostic Indium111-cG250 imaging confirmed cG250 accumulation in 23 patients with progressive ccRCC metastases (68). In this phase I study, patients received a high-activity dose of ¹⁷⁷Lu-cG250. At 3 months after the initial treatment, 74% of patients had stable illness (evaluated by PERCIST), and one patient had a partial response that persisted for 9 months. Mean growth of target tumor lesions was reduced from 40.4% in the 3 months before treatment to 5.5% 3 months after the first treatment cycle. There were no significant non-hematological adverse effects noted (68).

Additionally, in a phase II study, 14 patients with ccRCC were treated with ¹⁷⁷Lu-cG250, of which eight patients had stable disease and one had partial regression. The treatment was generally well tolerated (69).

Metastatic bone tumors and ¹⁷⁷Lu-EDTMP. A study by Elboga *et al* (70) found that ¹⁷⁷Lu-ethylenediamine tetramethyle nephosphonic acid (EDTMP) effectively relieved pain caused by bone metastases in patients with breast or prostate cancer. Of the 75 patients treated with ¹⁷⁷Lu-EDTMP, 59 responded positively, while 16 did not. The pain score was analyzed, and patients who responded had markedly lower pain scores after each radiopharmaceutical treatment. A meta-analysis involving 172 patients revealed that ¹⁷⁷Lu-EDTMP had a significant effect on relieving bone pain, suggesting that this agent could be a good choice when other pain-relieving radiopharmaceuticals are not available (71).

Currently, it is proposed that a ¹⁷⁷Lu-EDTMP rapid kit be developed based on its success in palliative treatment, which, if successfully implemented, could greatly reduce pain in terminal cancer patients (72).

6. Present problems and future research directions

At present, the main problem with ¹⁷⁷Lu TRT is the limited variety of drugs used in clinical practice, and the need for further improvement in clinical efficacy. ¹⁷⁷Lu TRT is a promising treatment method for solid tumors, but currently only ¹⁷⁷Lu-DOTA-TATE and ¹⁷⁷Lu-PSMA-617 have been approved treatment for NETs and prostate cancer, respectively. Moreover, the response rate of ¹⁷⁷Lu TRT for mCRPC was from 32.3 to 68%, and ¹⁷⁷Lu-DOTA-TATE for neuroendocrine tumors was only 18%.

Future research on ¹⁷⁷Lu TRT should focus on expanding the application of ¹⁷⁷Lu-DOTA-TATE and ¹⁷⁷Lu-PSMA-617 to other diseases that express PSMA or SSTR. There are currently ongoing clinical trials assessing the activity of ¹⁷⁷Lu-DOTA-TATE and ¹⁷⁷Lu-PSMA-617 on SSTR and PSMA-expressing tumors as listed in Table I. This includes

breast cancer, metastatic nasopharyngeal carcinoma, neuroblastoma, glioblastoma and glioma, NETs of the lung and thymus for SSTR-expressing tumors and upper metastatic gastric cancer for PSMA-expressing tumors.

Likewise, future research should focus on the development of more ¹⁷⁷Lu TRT targets in solid tumors. In the past 5 years, the ongoing clinical trials of new targets for ¹⁷⁷Lu TRT have included fibronectin ED-B, NY108, TLX591, LNC1004, 3PRGD2 and GD2 (Table II).

Lastly, the use of ¹⁷⁷Lu TRT in combination with other treatments should be explored. For example, in a preclinical model (murine model), ¹⁷⁷Lu-FAP-2287 enhanced anti-PD-1-mediated tumor growth inhibition by modulating the tumor microenvironment and increasing the recruitment of tumor-infiltrating CD8⁺ T cells (50). Another study showed that concurrent rather than sequential blockade of the PD-1/PD-L1 axis combined with ¹⁷⁷Lu TRT improves OS and long-term tumor control (73). The study of the metabolism principle of nuclide in the body can further clarify the treatment method in the future to improve the survival period. A Phase I study of the ¹⁷⁷Lu-DOTA⁰-Tyr³-Octreotate combination with nivolumab was well tolerated and showed signs of antitumor activity for patients with neuroendocrine tumors of the lung (74). In the past 5 years, the number of ongoing clinical trials of ¹⁷⁷Lu TRT combined with other treatments for solid tumors has increased. A list of the current trials is shown in Table III. The results of these clinical trials will likely improve the current options for ¹⁷⁷Lu TRT treatments.

7. Conclusions

¹⁷⁷Lu TRT have been successfully applied in patients with NETs or metastatic prostate cancer, resulting in prolonged PFS, OS and improved quality of life with tolerable toxicities. However, ¹⁷⁷Lu TRT are rarely used to treat other solid tumors and clinical efficacies need to be improved. The use of ¹⁷⁷Lu-DOTA-TATE and ¹⁷⁷Lu-PSMA-617 should be expanded to various diseases, and more new targets for ¹⁷⁷Lu TRT should be researched and developed. Lastly, ¹⁷⁷Lu TRT in combination with other treatments provide further treatment options for solid tumors.

Acknowledgements

Not applicable.

Funding

This work was financially supported by the NHC Key Laboratory of Nuclear Technology Medical Transformation, Mianyang Central Hospital (grant nos. 2022HYX001 and 2022HYX0015).

Availability of data and materials

Not applicable.

Authors' contributions

TN and MF drafted the manuscript. BL, FG and BT participated in the data review and collection for the study. XD

conceived the study and reviewed and edited the manuscript. All authors read and approved the final version of the manuscript. Data authentication is not applicable.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A and Bray F: global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 71: 209-249, 2021.
- Al-Toubah T, Strosberg J, Hallanger-Johnson J and El-Haddad G: Targeted radionuclide therapy in endocrine-related cancers: Advances in the last decade. *Front Endocrinol (Lausanne)* 14: 1187870, 2023.
- Salih S, Alkatheeri A, Alomaim W and Elliyanti A: Radiopharmaceutical treatments for cancer therapy, radionuclides characteristics, applications, and challenges. *Molecules* 27: 5231, 2022.
- Park UJ, Lee JS, Choi KH, Nam SS and Yu KH: Lu-177 preparation for radiotherapy application. *Appl Radiat Isot* 115: 8-12, 2016.
- Zakaly HMH, Mostafa MYA, Deryabina D and Zhukovsky M: Comparative studies on the potential use of ¹⁷⁷Lu-based radiopharmaceuticals for the palliative therapy of bone metastases. *Int J Radiat Biol* 96: 779-789, 2020.
- Dash A, Pillai MR and Knapp FF Jr: Production of (¹⁷⁷)Lu for targeted radionuclide therapy: Available options. *Nucl Med Mol Imaging* 49: 85-107, 2015.
- Tan HY, Yeong CH, Wong YH, McKenzie M, Kasbollah A, Shah MNM and Perkins AC: Neutron-activated theranostic radionuclides for nuclear medicine. *Nucl Med Biol* 90-91: 55-68, 2020.
- Cao J, Chen Y, Hu M and Zhang W: ¹⁷⁷Lu-PSMA-RLT of metastatic castration-resistant prostate cancer: Limitations and improvements. *Ann Nucl Med* 35: 861-870, 2021.
- Boegemann M, Schrader AJ and Rahbar K: ¹⁷⁷Lu-PSMA therapy: Current evidence for use in the treatment of patients with metastatic prostate cancer. *Urologe A* 56: 1440-1444, 2017 (In German).
- Ghosh A and Heston WD: Tumor target prostate specific membrane antigen (PSMA) and its regulation in prostate cancer. *J Cell Biochem* 91: 528-539, 2004.
- Fendler WP, Rahbar K, Herrmann K, Kratochwil C and Eiber M: ¹⁷⁷Lu-PSMA radioligand therapy for prostate cancer. *J Nucl Med* 58: 1196-1200, 2017.
- Sadaghiani MS, Sheikhabaei S, Werner RA, Pienta KJ, Pomper MG, Gorin MA, Solnes LB and Rowe SP: ¹⁷⁷Lu-PSMA radioligand therapy effectiveness in metastatic castration-resistant prostate cancer: An updated systematic review and meta-analysis. *Prostate* 82: 826-835, 2022.
- Weineisen M, Schottelius M, Simecek J, Baum RP, Yildiz A, Beykan S, Kulkarni HR, Lassmann M, Klette I, Eiber M, *et al*: 68Ga- and ¹⁷⁷Lu-labeled PSMA I&T: Optimization of a PSMA-targeted theranostic concept and first proof-of-concept human studies. *J Nucl Med* 56: 1169-1176, 2015.
- Barna S, Haug AR, Hartenbach M, Rasul S, Grubmüller B, Kramer G and Blaickner M: Dose calculations and dose-effect relationships in ¹⁷⁷Lu-PSMA I&T radionuclide therapy for metastatic castration-resistant prostate cancer. *Clin Nucl Med* 45: 661-667, 2020.
- Hartrampf PE, Weinzierl FX, Serfling SE, Pomper MG, Rowe SP, Higuchi T, Seitz AK, Kübler H, Buck AK and Werner RA: Hematotoxicity and nephrotoxicity in prostate cancer patients undergoing radioligand therapy with [¹⁷⁷Lu]Lu-PSMA I&T. *Cancers (Basel)* 14, 647, 2022.
- Bu T, Zhang L, Yu F, Yao X, Wu W, Zhang P, Shi L, Zang S, Meng Q, Ni Y, *et al*: ¹⁷⁷Lu-PSMA-I&T radioligand therapy for treating metastatic castration-resistant prostate cancer: A single-centre study in East Asians. *Front Oncol* 12: 835956, 2022.
- Emmett L, John N, Pathmanandavel S, Counter W, Ayers M, Sharma S, Agrawal S, Poole A, Hovey E, Pranavan G, *et al*: Patient outcomes following a response biomarker-guided approach to treatment using ¹⁷⁷Lu-PSMA-I&T in men with metastatic castrate-resistant prostate cancer (Re-SPECT). *Ther Adv Med Oncol* 15: 17588359231156392, 2023.
- Karimzadeh A, Heck M, Tauber R, Knorr K, Haller B, D'Alessandria C, Weber WA, Eiber M and Rauscher I: ¹⁷⁷Lu-PSMA-I&T for treatment of metastatic castration-resistant prostate cancer: Prognostic value of scintigraphic and clinical biomarkers. *J Nucl Med* 64: 402-409, 2023.
- Golan S, Frumer M, Zohar Y, Rosenbaum E, Yakimov M, Kedar D, Margel D, Baniel J, Steinmetz AP, Groshar D, *et al*: Neoadjuvant ¹⁷⁷Lu-PSMA-I&T radionuclide treatment in patients with high-risk prostate cancer before radical prostatectomy: A single-arm phase I trial. *Eur Urol Oncol* 6: 151-159, 2023.
- Stasiuk GJ and Long NJ: The ubiquitous DOTA and its derivatives: The impact of 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid on biomedical imaging. *Chem Commun (Camb)* 49: 2732-2746, 2013.
- Keam SJ: Lutetium Lu 177 vipivotide tetraxetan: First approval. *Mol Diagn Ther* 26: 467-475, 2022.
- Liu X, Fang GC, Lu H, Shi ZD, Chen ZS and Han CH: Lutetium Lu 177 vipivotide tetraxetan for prostate cancer. *Drugs Today (Barc)* 59: 37-49, 2023.
- Kim YJ and Kim YI: Therapeutic responses and survival effects of ¹⁷⁷Lu-PSMA-617 radioligand therapy in metastatic castrate-resistant prostate cancer: A meta-analysis. *Clin Nucl Med* 43: 728-734, 2018.
- Violet J, Sandhu S, Irvani A, Ferdinandus J, Thang SP, Kong G, Kumar AR, Akhurst T, Pattison DA, Beaulieu A, *et al*: Long-term follow-up and outcomes of retreatment in an expanded 50-patient single-center phase II prospective trial of ¹⁷⁷Lu-PSMA-617 theranostics in metastatic castration-resistant prostate cancer. *J Nucl Med* 61: 857-865, 2020.
- Hofman MS, Violet J, Hicks RJ and Sandhu S: [¹⁷⁷Lu]-PSMA-617 radionuclide therapy in patients with metastatic castration-resistant prostate cancer-Author's reply. *Lancet Oncol* 19: e373, 2018.
- Sartor O, de Bono J, Chi KN, Fizazi K, Herrmann K, Rahbar K, Tagawa ST, Nordquist LT, Vaishampayan N, El-Haddad G, *et al*: Lutetium-177-PSMA-617 for metastatic castration-resistant prostate cancer. *N Engl J Med* 385: 1091-1103, 2021.
- Schwartz LH, Litière S, de Vries E, Ford R, Gwyther S, Mandrekar S, Shankar L, Bogaerts J, Chen A, Dancey J, *et al*: RECIST 1.1-Update and clarification: From the RECIST committee. *Eur J Cancer* 62: 132-137, 2016.
- Hartrampf PE, Weinzierl FX, Buck AK, Rowe SP, Higuchi T, Seitz AK, Kübler H, Schirbel A, Essler M, Bundschuh RA and Werner RA: Matched-pair analysis of [¹⁷⁷Lu]Lu-PSMA I&T and [¹⁷⁷Lu]Lu-PSMA-617 in patients with metastatic castration-resistant prostate cancer. *Eur J Nucl Med Mol Imaging* 49: 3269-3276, 2022.
- Schuchardt C, Zhang J, Kulkarni HR, Chen X, Müller D and Baum RP: Prostate-specific membrane antigen radioligand therapy using ¹⁷⁷Lu-PSMA I&T and ¹⁷⁷Lu-PSMA-617 in patients with metastatic castration-resistant prostate cancer: Comparison of safety, biodistribution, and dosimetry. *J Nucl Med* 63: 1199-1207, 2022.
- Fizazi K, Herrmann K, Krause BJ, Rahbar K, Chi KN, Morris MJ, Sartor O, Tagawa ST, Kendi AT, Vogelzang N, *et al*: Health-related quality of life and pain outcomes with [¹⁷⁷Lu]Lu-PSMA-617 plus standard of care versus standard of care in patients with metastatic castration-resistant prostate cancer (VISION): A multicentre, open-label, randomised, phase 3 trial. *Lancet Oncol* 24: 597-610, 2023.
- Niaz MO, Sun M, Ramirez-Fort MK and Niaz MJ: Review of lutetium-177-labeled anti-prostate-specific membrane antigen monoclonal antibody j591 for the treatment of metastatic castration-resistant prostate cancer. *Cureus* 12: e7107, 2020.
- Liu H, Rajasekaran AK, Moy P, Xia Y, Kim S, Navarro V, Rahmati R and Bander NH: Constitutive and antibody-induced internalization of prostate-specific membrane antigen. *Cancer Res* 58: 4055-4060, 1998.

33. Batra JS, Niaz MJ, Whang YE, Sheikh A, Thomas C, Christos P, Vallabhajosula S, Jhanwar YS, Molina AM, Nanus DM, *et al.*: Phase I trial of docetaxel plus lutetium-177-labeled anti-prostate-specific membrane antigen monoclonal antibody J591 (¹⁷⁷Lu-J591) for metastatic castration-resistant prostate cancer. *Urol Oncol* 38: 848.e9-848.e16, 2020.
34. Shin D, Ha S, O JH, Rhew SA, Yoon CE, Kwon HJ, Moon HW, Park YH, Park SY, Park C, *et al.*: A single dose of novel PSMA-targeting radiopharmaceutical agent [¹⁷⁷Lu]ludotadipep for patients with metastatic castration-resistant prostate cancer: Phase I clinical trial. *Cancers (Basel)* 14: 6225, 2022.
35. Jacoba IM and Weber HC: Biomarkers in gastroenteropancreatic neuroendocrine neoplasms. *Curr Opin Endocrinol Diabetes Obes* 30: 175-180, 2023.
36. Fernandes CJ, Leung G, Eads JR and Katona BW: Gastroenteropancreatic neuroendocrine tumors. *Gastroenterol Clin North Am* 51: 625-647, 2022.
37. Liu M, Wei L, Liu W, Chen S, Guan M, Zhang Y, Guo Z, Liu R and Xie P: Trends in incidence and survival in patients with gastrointestinal neuroendocrine tumors: A SEER database analysis, 1977-2016. *Front Oncol* 13: 1079575, 2023.
38. Harris PE and Zheronsekov K: The evolution of PRRT for the treatment of neuroendocrine tumors; What comes next? *Front Endocrinol (Lausanne)* 13: 941832, 2022.
39. Theodoropoulou M and Stalla GK: Somatostatin receptors: From signaling to clinical practice. *Front Neuroendocrinol* 34: 228-252, 2013.
40. Ichikawa Y, Kobayashi N, Takano S, Kato I, Endo K and Inoue T: Neuroendocrine tumor theranostics. *Cancer Sci* 113: 1930-1938, 2022.
41. Pokuri VK, Fong MK and Iyer R: Octreotide and lanreotide in gastroenteropancreatic neuroendocrine tumors. *Curr Oncol Rep* 18: 7, 2016.
42. Das S, Al-Toubah T, El-Haddad G and Strosberg J: ¹⁷⁷Lu-DOTATATE for the treatment of gastroenteropancreatic neuroendocrine tumors. *Expert Rev Gastroenterol Hepatol* 13: 1023-1031, 2019.
43. Strosberg J, El-Haddad G, Wolin E, Hendifar A, Yao J, Chasen B, Mittra E, Kunz PL, Kulke MH, Jacene H, *et al.*: Phase 3 trial of ¹⁷⁷Lu-Dotatate for midgut neuroendocrine tumors. *N Engl J Med* 376: 125-135, 2017.
44. Strosberg JR, Caplin ME, Kunz PL, Ruzsniwski PB, Bodei L, Hendifar A, Mittra E, Wolin EM, Yao JC, Pavel ME, *et al.*: ¹⁷⁷Lu-Dotatate plus long-acting octreotide versus high-dose long-acting octreotide in patients with midgut neuroendocrine tumours (NETTER-1): Final overall survival and long-term safety results from an open-label, randomised, controlled, phase 3 trial. *Lancet Oncol* 22: 1752-1763, 2021.
45. Khan MS, Stamp E, Sammon C, Brabander T, de Herder WW and Pavel ME: Matching-adjusted indirect treatment comparison of [¹⁷⁷Lu]Lu-DOTA-TATE, everolimus and sunitinib in advanced, unresectable gastroenteropancreatic neuroendocrine tumours: Relative effectiveness of [¹⁷⁷Lu]Lu-DOTA-TATE in gastroenteropancreatic neuroendocrine tumours. *EJC Suppl* 16: 5-13, 2021.
46. Alsadik S, Gnanasegaran G, Chen L, Quigley AM, Mandair D, Toumpanakis C, Caplin M and Navalkissoor S: Safety and efficacy of ¹⁷⁷Lu-DOTATATE in neuroendocrine tumor patients with extensive bone disease. *Clin Nucl Med* 48: 667-672, 2023.
47. Albrecht J, Exner S, Grötzinger C, Prasad S, Konietschke F, Beindorff N, Kühl AA, Prasad V, Brenner W and Koziol EJ: Multimodal imaging of 2-cycle PRRT with ¹⁷⁷Lu-DOTA-JR11 and ¹⁷⁷Lu-DOTATOC in an orthotopic neuroendocrine xenograft tumor mouse model. *J Nucl Med* 62: 393-398, 2021.
48. Mansi R, Plas P, Vauquelin G and Fani M: Distinct in vitro binding profile of the somatostatin receptor subtype 2 antagonist [¹⁷⁷Lu]Lu-OPS201 compared to the agonist [¹⁷⁷Lu]Lu-DOTA-TATE. *Pharmaceuticals (Basel)* 14: 1265, 2021.
49. Zboralski D, Hoehne A, Bredenbeck A, Schumann A, Nguyen M, Schneider E, Ungewiss J, Paschke M, Haase C, von Hacht JL, *et al.*: Preclinical evaluation of FAP-2286 for fibroblast activation protein targeted radionuclide imaging and therapy. *Eur J Nucl Med Mol Imaging* 49: 3651-3667, 2022.
50. Zboralski D, Osterkamp F, Christensen E, Bredenbeck A, Schumann A, Hoehne A, Schneider E, Paschke M, Ungewiss J, Haase C, *et al.*: Fibroblast activation protein targeted radiotherapy induces an immunogenic tumor microenvironment and enhances the efficacy of PD-1 immune checkpoint inhibition. *Eur J Nucl Med Mol Imaging* 50: 2621-2635, 2023.
51. Baum RP, Schuchardt C, Singh A, Chantadisai M, Robiller FC, Zhang J, Mueller D, Eismant A, Almaguel F, Zboralski D, *et al.*: Feasibility, biodistribution, and preliminary dosimetry in peptide-targeted radionuclide therapy of diverse adenocarcinomas using ¹⁷⁷Lu-FAP-2286: First-in-humans results. *J Nucl Med* 63: 415-423, 2022.
52. Minczeles NS, Bos EM, de Leeuw RC, Kros JM, Konijnenberg MW, Bromberg JEC, de Herder WW, Dirven CMF, Hofland J and Brabander T: Efficacy and safety of peptide receptor radionuclide therapy with [¹⁷⁷Lu]Lu-DOTA-TATE in 15 patients with progressive treatment-refractory meningioma. *Eur J Nucl Med Mol Imaging* 50: 1195-1204, 2023.
53. Nishida H, Kondo Y, Kusaba T, Kadowaki H and Daa T: Immunohistochemical reactivity of prostate-specific membrane antigen in salivary gland tumors. *Head Neck Pathol* 16: 427-433, 2022.
54. van Boxtel W, Uijen MJM, Verhaegh GW, Willems SM, Jonker MA; PALGA Group; Schalken JA, van Engen-van Grunsven ICH and van Herpen CML: Prognostic value of PSMA, c-MET and E-cadherin in salivary duct carcinoma. *Oral Oncol* 110: 105018, 2020.
55. van Boxtel W, Lütje S, van Engen-van Grunsven ICH, Verhaegh GW, Schalken JA, Jonker MA, Nagarajah J, Gotthardt M and van Herpen CML: ⁶⁸Ga-PSMA-HBED-CC PET/CT imaging for adenoid cystic carcinoma and salivary duct carcinoma: A phase 2 imaging study. *Theranostics* 10: 2273-2283, 2020.
56. Uijen MJM, Derks YHW, Merckx RIJ, Schilham MGM, Roosen J, Privé BM, van Lith SAM, van Herpen CML, Gotthardt M, Heskamp S, *et al.*: PSMA radioligand therapy for solid tumors other than prostate cancer: Background, opportunities, challenges, and first clinical reports. *Eur J Nucl Med Mol Imaging* 48: 4350-4368, 2021.
57. Lütje S, Sauerwein W, Lauenstein T, Bockisch A and Poeppel TD: In vivo visualization of prostate-specific membrane antigen in adenoid cystic carcinoma of the salivary gland. *Clin Nucl Med* 41: 476-477, 2016.
58. Klein Nulent TJW, van Es RJJ, Willems SM, Braat AJAT, Devriese LA, de Bree R and de Keizer B: First experiences with ¹⁷⁷Lu-PSMA-617 therapy for recurrent or metastatic salivary gland cancer. *EJNMMI Res* 11: 126, 2021.
59. Terroir M, Lamesa C, Krim M, Vija L, Texier JS, Cassou-Mounat T, Delord JP, Vallot D and Courbon F: Radioligand therapy with [¹⁷⁷Lu]Lu-PSMA-617 for salivary gland cancers: Literature review and first compassionate use in France. *Pharmaceuticals (Basel)* 16: 754, 2023.
60. Lütje S, Gomez B, Cohnen J, Umutlu L, Gotthardt M, Poeppel TD, Bockisch A and Rosenbaum-Krumme S: Imaging of prostate-specific membrane antigen expression in metastatic differentiated thyroid cancer using ⁶⁸Ga-HBED-CC-PSMA PET/CT. *Clin Nucl Med* 42: 20-25, 2017.
61. Lawhn-Heath C, Yom SS, Liu C, Villanueva-Meyer JE, Aslam M, Smith R, Narwal M, Juarez R, Behr SC, Pampaloni MH, *et al.*: Gallium-68 prostate-specific membrane antigen (⁶⁸Ga]Ga-PSMA-11) PET for imaging of thyroid cancer: A feasibility study. *EJNMMI Res* 10: 128, 2020.
62. Sollini M, Kirienko M, di Tommaso L, Pini C, Gelardi F, Ariano S, Lania AG, Mazziotti G, Mercante G and Chiti A: The complementary role of PSMA expression and [¹⁸F]FDG PET/CT in predicting thyroid cancer outcome: From black and white to shades of gray, in the era of precision oncology. *EJNMMI Res* 13: 54, 2023.
63. Pitalua-Cortes Q, García-Perez FO, Vargas-Ahumada J, Gonzalez-Rueda S, Gomez-Argumosa E, Ignacio-Alvarez E, Soldevilla-Gallardo I and Torres-Agredo L: Head-to-head comparison of ⁶⁸Ga-PSMA-11 and ¹³¹I in the follow-up of well-differentiated metastatic thyroid cancer: A new potential theragnostic agent. *Front Endocrinol (Lausanne)* 12: 794759, 2021.
64. Assadi M and Ahmadzadehfah H: ¹⁷⁷Lu-DOTATATE and ¹⁷⁷Lu-prostate-specific membrane antigen therapy in a patient with advanced metastatic radioiodine-refractory differentiated thyroid cancer after failure of tyrosine kinase inhibitors treatment. *World J Nucl Med* 18: 406-408, 2019.
65. de Vries LH, Lodewijk L, Braat AJAT, Krijger GC, Valk GD, Lam MGEH, Borel Rinkes IHM, Vriens MR and de Keizer B: ⁶⁸Ga-PSMA PET/CT in radioactive iodine-refractory differentiated thyroid cancer and first treatment results with ¹⁷⁷Lu-PSMA-617. *EJNMMI Res* 10: 18, 2020.

66. Pastoreková S, Parkkila S, Parkkila AK, Opavský R, Zelník V, Saarnio J and Pastorek J: Carbonic anhydrase IX, MN/CA IX: Analysis of stomach complementary DNA sequence and expression in human and rat alimentary tracts. *Gastroenterology* 112: 398-408, 1997.
67. Steffens MG, Oosterwijk-Wakka JC, Zegwaard-Hagemeier NE, Boerman OC, Debruyne FM, Corstens FH and Oosterwijk E: Immunohistochemical analysis of tumor antigen saturation following injection of monoclonal antibody G250. *Anticancer Res* 19: 1197-1200, 1999.
68. Stillebroer AB, Boerman OC, Desar IME, Boers-Sonderen MJ, van Herpen CML, Langenhuijsen JF, Smith-Jones PM, Oosterwijk E, Oyen WJ and Mulders PF: Phase I radioimmunotherapy study with lutetium 177-labeled anti-carbonic anhydrase IX monoclonal antibody girentuximab in patients with advanced renal cell carcinoma. *Eur Urol* 64: 478-485, 2013.
69. Muselaers CH, Boers-Sonderen MJ, van Oostenbrugge TJ, Boerman OC, Desar IM, Stillebroer AB, Mulder SF, van Herpen CM, Langenhuijsen JF, Oosterwijk E, *et al*: Phase 2 study of lutetium 177-labeled anti-carbonic anhydrase IX monoclonal antibody girentuximab in patients with advanced renal cell carcinoma. *Eur Urol* 69: 767-770, 2016.
70. Elboga U, Kilbas B, Sahin E, Cayırlı YB, Eryılmaz K, Begec T, Bakar HE, Mercanoglu G and Celen YZ: An automated synthesis of ¹⁷⁷Lu-EDTMP as an efficient bone-seeking therapeutic radiopharmaceutical. *Eur Rev Med Pharmacol Sci* 25: 4829-4834, 2021.
71. Askari E, Harsini S, Vahidfar N, Divband G and Sadeghi R: ¹⁷⁷Lu-EDTMP for metastatic bone pain palliation: A systematic review and meta-analysis. *Cancer Biother Radiopharm* 36: 383-390, 2021.
72. Mercanoglu G, Zilbeyaz K and Arslan N: Synthesis and ready to use kit formulation of EDTMP for the preparation of ¹⁷⁷Lu-EDTMP as a bone palliation radiopharmaceutical. *Curr Radiopharm* 16: 38-43, 2023.
73. Chen H, Zhao L, Fu K, Lin Q, Wen X, Jacobson O, Sun L, Wu H, Zhang X, Guo Z, *et al*: Integrin $\alpha_3\beta_3$ -targeted radionuclide therapy combined with immune checkpoint blockade immunotherapy synergistically enhances anti-tumor efficacy. *Theranostics* 9: 7948-7960, 2019.
74. Kim C, Liu SV, Subramaniam DS, Torres T, Loda M, Esposito G and Giaccone G: Phase I study of the ¹⁷⁷Lu-DOTA⁰-Tyr³-octreotate (lutathera) in combination with nivolumab in patients with neuroendocrine tumors of the lung. *J Immunother Cancer* 8: e000980, 2020.



Copyright © 2024 Niu et al. This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.